

New Heavy-Light Mesons $Q\bar{q}$

Takayuki Matsuki*

Tokyo Kasei University, 1-18-1 Kaga, Itabashi, Tokyo 173-8602, JAPAN

Toshiyuki Morii†

*Graduate School of Science and Technology, Kobe University,
Nada, Kobe 657-8501, JAPAN*

Kazutaka Sudoh‡

*Institute of Particle and Nuclear Studies,
High Energy Accelerator Research Organization,
1-1 Oho, Tsukuba, Ibaraki 305-0801, JAPAN*

(Dated: February 2, 2008)

We succeed in reproducing the $\ell = 1$ B mesons, $B_1(5720)$, $B_2^*(5745)$, and $B_{s2}^*(5839)$ that are recently reported by D0 and CDF, by using our semirelativistic quark potential model, which also succeeded in predicting the mass spectra of the narrow D_{sJ} as well as broad $D_0^*(0^+)$ and $D_1'(1^+)$ particles a couple of years ago.

Mass of higher excited states ($\ell = 1, 2$) of B and B_s mesons, which are not yet observed, is also predicted at the first order in p/m_b with internal quark momentum p and the b quark mass m_b . We find the corresponding B_{sJ} are below BK/B^*K threshold and should have narrow decay widths contrary to most other predictions. Also already established states ($\ell = 0$ and $\ell = 1$) of D , D_s , B , and B_s heavy mesons are simultaneously reproduced in good agreement with experimental data within one percent of accuracy. To calculate these D/D_s and B/B_s heavy mesons we use different values of strong coupling α_s .

PACS numbers: 12.39.Hg, 12.39.Pn, 12.40.Yx, 14.40.Lb, 14.40.Nd

Keywords: potential model; spectroscopy; heavy mesons

I. INTRODUCTION

Recently discovered were narrow meson states, $D_{s0}(2317)$ by BaBar [1] and $D_{s1}(2460)$ by CLEO [2], both of which were confirmed by Belle [3]. These are identified as $j^P = 0^+$ and 1^+ of $c\bar{s}$ excited ($\ell = 1$) bound states, respectively, and have caused a revival of study on heavy meson spectroscopy. Subsequently, another set of broad heavy mesons, $D_0^*(2308)$ and $D_1'(2427)$ were discovered by Belle collaboration [4]. Those are identified as $c\bar{q}$ ($q = u/d$) excited ($\ell = 1$) bound states and have the same quantum numbers $j^P = 0^+$ and 1^+ as D_{sJ} , respectively. The decay widths of these excited D_{sJ} mesons are narrow since the masses are below DK/D^*K threshold and hence the dominant decay modes violate the isospin invariance, whereas those excited D mesons are broad because of no such restriction as in D_{sJ} cases. More recent experiments reported by CDF and D0 [5] found narrow B and B_s states of $\ell = 1$, $B_1(5720)$, $B_2^*(5745)$, and $B_{s2}^*(5839)$. These are narrow because these decay through the D-waves.

There are some discussions [6] that a quark potential model is not appropriate to describe these new states. In fact, mass spectra and decay widths of these states could not be reproduced by relativised/relativistic quark potential models [7, 8, 9]. In [7, 8], for instance, kinetic terms of quarks are taken as positive definite $\sqrt{p^2 + m^2}$ and spin-independent linear-rising confining and short-range Coulomb potentials are taken into account together with spin-dependent interaction terms symmetric in Q and $\bar{q}$. In [9], even though degeneracy among members of a spin doublet in the limit of heavy quark symmetry is taken into account, they obtained masses more than a hundred MeV larger than the observed D_{sJ} mesons. The nonrelativistic quark potential model applied both to mesons and baryons has been so successful in reproducing low lying hadrons (see the review [10]) that people were puzzled why it did not work for D_{sJ} mesons (atom like mesons).

To understand these heavy meson states, a number of interesting ideas have been proposed so far. One is an effective Lagrangian approach. The authors of [11, 12] have proposed the modified Goldberger-Treiman relation in an effective

*E-mail: matsuki@tokyo-kasei.ac.jp

†E-mail: morii@kobe-u.ac.jp

‡E-mail: kazutaka.sudoh@kek.jp

Lagrangian, by which the mass difference between two spin doublets ($0^-, 1^-$) and ($0^+, 1^+$) are reproduced. This idea is theoretically successful in explaining D_s states but in fail for D states though experimentally the relation holds. Other approaches proposed to understand these new states are the QCD sum rule [13], implications of a two-meson molecule [6, 14], four quark system [15, 16, 17], lattice calculation [18] and so on. The Bethe-Salpeter (BS) equation can be an alternative way to take relativistic corrections into account.[19, 20, 21] A coupled channel method [22] is another candidate to interpret the new heavy mesons though underlying physics is unclear. These approaches are not yet completely established even though they may be interesting themselves. Hence, the problem still remains unsolved by these approaches and is challenging. See the detailed review [23] on heavy mesons.

We believe that a quark potential model still powerfully survives if we treat a bound state equation appropriately, being able to predict not only mass differences but also absolute values of hadron masses. In fact, long time ago before the discovery of $D_{s0}(2317)$ by BaBar, two of us (T.M. and T.M.) [24] proposed a new bound state equation for atom-like mesons, i.e., heavy mesons composed of a heavy quark Q and a light antiquark $\bar{q}$, in which quarks are treated as four-spinor Dirac particles, and Hamiltonian, wave functions, and eigenvalues are all expanded in p/m_Q so that a light quark is treated as relativistically as possible and nonrelativistic reduction is made for a heavy quark. The heavy quark symmetry is taken into account consistently within a potential model. It is remarkable that the model could predict the levels of $D_{s0}(2317)$, $D_{s1}(2460)$, $D_0^*(2308)$, and $D_1'(2427)$ quite successfully even using rather obscure parameter values due to a small number of data (ground states and only a few excited states) at the time of publication [24].

Recent experiments on B and B_s (discovery of states of $\ell = 1$, $B_1(5720)$, $B_2^*(5745)$, and $B_{s2}^*(5839)$) by CDF and D0 [5] serves us a good testing ground whether our formulation, semirelativistic quark potential model, or other formulation also works or not. One of the main purposes of this paper is to show that our model can also reproduce these B/B_s states within one percent of accuracy compared with the experiments. Another aim of this paper is to reproduce and predict the whole spectrum of D/D_s and B/B_s heavy mesons, including higher states, 3D_1 and $^3D_2/1D_2$, which we failed to have reasonable values in [24].

In the previous paper [24], because of the small number of input data in those days, we searched for a *local minimum* which fits with at least the lowest ground states and found that the χ^2 analysis led to a *negative* optimal value of b in a scalar potential $S(r) = b + r/a^2$. However, in this paper we have obtained a *positive* b owing to enough experimental input data this time using the Minuit. One difference caused by this sign change appears in light quark mass, $m_q = m_u = m_d$, whose effective mass in the lowest order Hamiltonian becomes $m_q + b$. This effective mass becomes negative, ~ -38 MeV for $b < 0$ in [24] and positive, ~ 86 MeV in this paper. Heavy quark masses, m_c and m_b , are also affected by this sign change, i.e., we have obtained a few hundred MeV smaller values in this paper than those in [24]. Other significant influence of sign change is on the magnitude of the corrections of lower components of a heavy quark, which are comparable to upper component contributions in [24] ($b < 0$), while in this paper the lower component contributions are comparable to the second order contributions in p/m_Q as we will see later. These are the main difference from [24] in the way of data analysis and otherwise we are using the same approach as before. In addition to a good result for mass levels of $\ell = 0$ and $\ell = 1$ states of all D , D_s , B and B_s mesons, a positive b causes not only obtaining plausible values for $k = 2$ ($\ell = 2$) states (corresponding to 3D_1 , $^3D_2/1D_2$) but also obtaining relatively small degenerate mass M_0 in a spin multiplet so that the corrections become larger in this paper compared with [24]. That is, the sign of b is important to obtain the whole spectra of heavy mesons even though the lower lying spectra are correctly predicted in both cases. Another important and different point from the former paper is that we treat even the light quark masses (m_u , m_d , and m_s) as free parameters in this paper, while they were given as input in [24].

In this paper, using the refined parameters obtained from recent new experimental data, we elaborate computation of the mass spectra of heavy mesons D , D_s , B , and B_s mesons, including $D_{s0}(2317)$, $D_{s1}(2460)$, $D_0^*(2308)$, $D_1'(2427)$, $B_1(5720)$, $B_2^*(5745)$, and $B_{s2}^*(5839)$. We find that in this new analysis all of the masses of not only above mentioned new excited ($\ell = 1$) states $j^P = 0^+, 1^+$, and 2^+ but also those of other excited ($\ell = 1$) states $j^P = 1^+$ and 2^+ , and ground ($\ell = 0$) states $j^P = 0^-$ and 1^- of D , D_s , B , and B_s mesons can be reproduced well within one percent of accuracy. Furthermore, we *predict* B_{s0} and B_{s1} are below BK/B^*K threshold, which means one can find them as narrow states and is the same results as in [25] although we have about 100 MeV smaller values for 0^+ and 1^+ states. This is contrary to most other quark potential model predictions as in D_{sJ} particles.

The main difference between our treatment of the quark potential model and others is that we have systematically expanded interaction terms in p/m_Q and that we have taken into account lower components of a heavy quark in a bound state. It turns out that *systematically expanded interaction terms are quite different from those of nonrelativistic and/or relativised models* except for [21]. In [21] they used the BS equation and projected heavy quark sector on positive states, otherwise their final formulation is very close to ours. Another large difference is the magnitude of a light quark mass, i.e., other models including [21] use a constituent quark mass while we got much smaller values, e.g., $m_u, m_d \sim 10$ MeV and $m_s \sim 100$ MeV.

The paper is organized as follows. In the next section, we briefly review our model, where some necessary formulas

are presented for readers' understanding and a discussion on a convenient quantum number k is given. In Sec. III, numerical results are presented in Tables III and IV for the mass levels of D and D_s mesons including their excited states together with the most optimal values of parameters in Table II, where we find that calculated masses are in good agreement with all existing experimental data. Those of B and B_s mesons are also given in the same section, Tables VI and VII. To show smallness of the second order corrections, we calculate D meson mass spectra with the second order corrections and give them in Table V. Section IV is devoted to conclusion and discussion, including the reason why our semirelativistic quark potential model approach gives good results compared with others.

II. BRIEF REVIEW OF OUR SEMIRELATIVISTIC MODEL

Let us start with an effective two-body Hamiltonian for a system of a heavy quark Q with mass m_Q and a light antiquark $\bar{q}$ with mass $m_{\bar{q}}$ [26, 27],¹

$$H = (\vec{\alpha}_{\bar{q}} \cdot \vec{p}_{\bar{q}} + \beta_{\bar{q}} m_{\bar{q}}) + (\vec{\alpha}_Q \cdot \vec{p}_Q + \beta_Q m_Q) + \beta_{\bar{q}} \beta_Q S(r) + \left\{ 1 - \frac{1}{2} [\vec{\alpha}_{\bar{q}} \cdot \vec{\alpha}_Q + (\vec{\alpha}_{\bar{q}} \cdot \vec{n})(\vec{\alpha}_Q \cdot \vec{n})] \right\} V(r), \quad (1)$$

where S and V are a scalar confining potential and a vector one-gluon exchange Coulomb potential with transverse interaction and $\vec{n} = \vec{r}/r$ is a unit vector along the relative coordinate between Q and $\bar{q}$. S and V are given explicitly as

$$S(r) = \frac{r}{a^2} + b, \quad V(r) = -\frac{4}{3} \frac{\alpha_s}{r}, \quad (2)$$

where a and $1/b$ are parameters with length dimension and α_s is a strong coupling constant. Since a heavy quark Q is sufficiently heavier than a light antiquark $\bar{q}$ which is orbiting Q , it is reasonable to apply the Foldy–Wouthuysen–Tani (FWT) transformation [29] to the heavy quark related operators in H of Eq.(1). One should notice that the FWT is introduced so that free kinetic terms of a heavy quark become $\beta_Q m_Q$ and one can expand the interaction terms in terms of p/m_Q systematically,² which is equivalent to a nonrelativistic reduction of the original model. By doing this way, we can get an eigenvalue equation for an atom-like bound state $Q\bar{q}$ [24]³

$$(H_{\text{FWT}} - m_Q) \otimes \psi_{\text{FWT}} = \tilde{E} \psi_{\text{FWT}}, \quad (3)$$

with

$$H_{\text{FWT}} - m_Q = H_{-1} + H_0 + H_1 + H_2 + \cdots, \quad (4)$$

where $\tilde{E} = E - m_Q$ (E being the bound state mass of $Q\bar{q}$) is the binding energy and a notation $\otimes$ denotes that gamma matrices of a light antiquark is multiplied from left with the wave function while those of a heavy quark from right. H_i in Eq.(4) denotes the i -th order expanded Hamiltonian in p/m_Q and its explicit expressions are presented in [24] and also in the Appendix A. Then, Eq.(3) can be numerically solved in order by order in p/m_Q according to the standard perturbation method by expanding the eigenvalue and the wave function as

$$\tilde{E} = E - m_Q = E_0^a + E_1^a + E_2^a + \cdots, \quad (5)$$

$$\psi_{\text{FWT}} = \psi_0^a + \psi_1^a + \psi_2^a + \cdots, \quad (6)$$

where a superscript a represents a set of quantum numbers, a total angular momentum j , its z component m and a quantum number k of the spinor operator K defined below in Eq.(11) and a subscript i of E_i^a and ψ_i^a stands for the i -th order in p/m_Q .

In practice, it is reasonable to first solve the 0-th order non-trivial equation by variation to get the 0-th order eigenvalues and their wave functions and then to estimate the corrections perturbatively in order by order in p/m_Q

¹ Miyazawa and Tanaka [28] pointed out that a relativistic bound state equation having this effective Hamiltonian can be derived with a boundary condition in relative time that is different from that of the Bethe–Salpeter equation.

² There is another way of expanding the Hamiltonian. See [30] where the Bloch method [31] is used to expand the effective Hamiltonian in p/m_Q , which is reproduced in Appendix B.

³ There is a pioneering work of this approach [32], where expanded terms of the Hamiltonian in p/m_Q are treated differently from ours.

by evaluating the matrix elements for those terms. The 0-th order eigenvalue E_0^a and the wave function ψ_0^a for the $Q\bar{q}$ bound state is obtained by solving the 0-th order eigenvalue equation,

$$H_0 \otimes \psi_0^a = E_0^a \psi_0^a, \quad H_0 = \vec{\alpha}_{\bar{q}} \cdot \vec{p}_{\bar{q}} + \beta_{\bar{q}} (m_{\bar{q}} + S(r)) + V(r). \quad (7)$$

The wave function which has two spinor indexes and is expressed as 4×4 matrix form is explicitly described by

$$\psi_0^a = \begin{pmatrix} 0 & \Psi_{jm}^k(\vec{r}) \end{pmatrix}, \quad (8)$$

$$\Psi_{jm}^k(\vec{r}) = \frac{1}{r} \begin{pmatrix} u_k(r) y_{jm}^k \\ iv_k(r) y_{jm}^{-k} \end{pmatrix}, \quad (9)$$

where y_{jm}^k being of 2×2 form are the angular part of the wave functions and $u_k(r)$ and $v_k(r)$ are the radial parts. The total angular momentum of a heavy meson $\vec{J}$ is the sum of the total angular momentum of the light quark degrees of freedom $\vec{S}_\ell$ and the heavy quark spin $\frac{1}{2}\vec{\Sigma}_Q$:

$$\vec{J} = \vec{S}_\ell + \frac{1}{2}\vec{\Sigma}_Q \quad \text{with} \quad \vec{S}_\ell = \vec{L} + \frac{1}{2}\vec{\Sigma}_{\bar{q}}, \quad (10)$$

where $\frac{1}{2}\vec{\Sigma}_{\bar{q}}$ ($= \frac{1}{2}\vec{\sigma}_{\bar{q}} \ 1_{2 \times 2}$) and $\vec{L}$ are the 4×4 spin and the orbital angular momentum of a light antiquark, respectively. Furthermore, k is the quantum number of the following spinor operator K [24]

$$K = -\beta_{\bar{q}} (\vec{\Sigma}_{\bar{q}} \cdot \vec{L} + 1), \quad K \Psi_{jm}^k = k \Psi_{jm}^k. \quad (11)$$

Note that in our approach K is introduced for a two-body bound system of $Q\bar{q}$, though the same form of the operator K can be defined in the case of a single Dirac particle in a central potential [33]. One can easily show [34] that the operator K^2 is equivalent to $\vec{S}_\ell^2$ and it holds $k = \pm (s_\ell + \frac{1}{2})$ or $s_\ell = |k| - \frac{1}{2}$. k is also related to j as

$$j = |k| \quad \text{or} \quad |k| - 1 \quad (k \neq 0). \quad (12)$$

In order to diagonalize the 0-th order Hamiltonian H_0 of Eq.(7) in the k representation, it is convenient to introduce a spinor representation of y_{jm}^k which is given by the unitary transformation (Eq.(14) below) of a set of the spherical harmonics Y_j^m and the vector spherical harmonics defined by

$$\vec{Y}_{jm}^{(L)} = -\vec{n} Y_j^m, \quad \vec{Y}_{jm}^{(E)} = \frac{r}{\sqrt{j(j+1)}} \vec{\nabla} Y_j^m, \quad \vec{Y}_{jm}^{(M)} = -i\vec{n} \times \vec{Y}_{jm}^{(E)}, \quad (13)$$

where $\vec{n} = \vec{r}/r$. These vector spherical harmonics are nothing but a set of eigenfunctions for a spin-1 particle. $\vec{Y}_{jm}^{(A)}$ ($A=L, M, E$) are eigenfunctions of $\vec{J}^2$ and J_z , having the eigenvalues $j(j+1)$ and m . The parity of these functions is assigned as $(-1)^{j+1}$, $(-1)^j$, $(-1)^{j+1}$ for $A=L, M$, and E , respectively, since Y_j^m has a parity $(-1)^j$. The unitary transformation is given by

$$\begin{pmatrix} y_{jm}^{-(j+1)} \\ y_{jm}^j \end{pmatrix} = U \begin{pmatrix} Y_j^m \\ \vec{\sigma} \cdot \vec{Y}_{jm}^{(M)} \end{pmatrix}, \quad \begin{pmatrix} y_{jm}^{j+1} \\ y_{jm}^{-j} \end{pmatrix} = U \begin{pmatrix} \vec{\sigma} \cdot \vec{Y}_{jm}^{(L)} \\ \vec{\sigma} \cdot \vec{Y}_{jm}^{(E)} \end{pmatrix}, \quad (14)$$

where the matrix U is defined by

$$U = \frac{1}{\sqrt{2j+1}} \begin{pmatrix} \sqrt{j+1} & \sqrt{j} \\ -\sqrt{j} & \sqrt{j+1} \end{pmatrix}. \quad (15)$$

Now, substituting Eqs.(8) and (9) into Eq.(7), one can eliminate the angular part of the wave function, y_{jm}^k , from the eigenvalue equation and obtain the radial equation as follows,

$$\begin{pmatrix} m_q + S + V & -\partial_r + \frac{k}{r} \\ \partial_r + \frac{k}{r} & -m_q - S + V \end{pmatrix} \begin{pmatrix} u_k(r) \\ v_k(r) \end{pmatrix} = E_0^k \begin{pmatrix} u_k(r) \\ v_k(r) \end{pmatrix}, \quad (16)$$

which depends on the quantum number k alone. This is just the same form as a one-body Dirac equation in a central potential. Since K commutes with H_0 and the states Ψ_{jm}^k have the same energy E_0^k for different values of j , these states having a total angular momentum $j = |k| - 1$ and $|k|$, a spin doublet, are degenerate with the same value of k .

TABLE I: States classified by various quantum numbers

j^P	0^-	1^-	0^+	1^+	1^+	2^+	1^-	2^-
k	-1	-1	1	1	-2	-2	2	2
$s_\ell^{\pi_\ell}$	$\frac{1}{2}^-$	$\frac{1}{2}^-$	$\frac{1}{2}^+$	$\frac{1}{2}^+$	$\frac{3}{2}^+$	$\frac{3}{2}^+$	$\frac{3}{2}^-$	$\frac{3}{2}^-$
$2s+1 l_j$	1S_0	3S_1	3P_0	$^3P_1, ^1P_1$	$^1P_1, ^3P_1$	3P_2	3D_1	$^3D_2, ^1D_2$
Ψ_j^k	Ψ_0^{-1}	Ψ_1^{-1}	Ψ_0^1	Ψ_1^1	Ψ_1^{-2}	Ψ_2^{-2}	Ψ_1^2	Ψ_2^2




FIG. 1: Procedure how the degeneracy is resolved in our model.

The parity of the eigenfunction ψ_0^a is defined by the upper (“large”) component of Eq.(9) and hence taking into account the intrinsic parity of quark and antiquark, the parity of the meson $Q\bar{q}$ is given by [34]

$$P = \frac{k}{|k|}(-1)^{|k|+1}, \quad (17)$$

which is equal to the parity π_ℓ of the light degrees of freedom in the heavy quark effective theory (HQET) as shown in Table I. It is remarkable that both the total angular momentum j and the parity P of a heavy meson can be simultaneously determined by k alone. Or the states can be completely classified in terms of k and j as shown in Table I.

Classification by the quantum number k is convenient to discuss the relation between the symmetry of the system and the structure of mass levels. Now, let us look at how masses of $Q\bar{q}$ mesons are generated in our model. First, when the light quark mass $m_{\bar{q}}$ and the scalar potential $S(r)$ are neglected, the 0-th order Hamiltonian H_0 has both a chiral symmetry and a heavy quark symmetry and the eigenvalues with the same value of $|k|$ are degenerate in this limit, where all masses of $0^-, 1^-, 0^+$, and 1^+ states with $|k| = 1$ are degenerate, as $m(0^-) = m(1^-) = m(0^+) = m(1^+) = m_Q$. [25] Then, by turning on the light quark mass and the scalar potential, $m_{\bar{q}} \neq 0$ and $S(r) \neq 0$, the chiral symmetry is broken and the degeneracy between two spin doublets ($0^-, 1^-$) and ($0^+, 1^+$) is resolved. At this stage, members in spin multiplets are still degenerate due to the same quantum number k . Finally, by adding p/m_Q corrections, the heavy quark symmetry is broken and the hyperfine splitting occurs as shown in Fig. 1.

III. NUMERICAL CALCULATION

To obtain masses of $Q\bar{q}$ states, we first solve the eigenvalue equation Eq.(16) by variation to get the lowest eigenvalues and the wave functions. By taking account of the asymptotic behavior at both $r \rightarrow 0$ and $r \rightarrow \infty$, the trial functions for $u_k(r)$ and $v_k(r)$ are given by

$$u_k(r), v_k(r) \sim w_k(r) \left(\frac{r}{a}\right)^\lambda \exp \left[-(m_q + b)r - \frac{1}{2} \left(\frac{r}{a}\right)^2 \right], \quad (18)$$

where

$$\lambda = \sqrt{k^2 - \left(\frac{4\alpha_s}{3}\right)^2}, \quad (19)$$

and $w_k(r)$ is a finite series of a polynomial in r

$$w_k(r) = \sum_{i=0}^{N-1} a_i^k \left(\frac{r}{a}\right)^i, \quad (20)$$

TABLE II: Most optimal values of parameters.

Parameters	α_s^c	α_s^b	a (GeV $^{-1}$)	b (GeV)	$m_{u,d}$ (GeV)	m_s (GeV)	m_c (GeV)	m_b (GeV)
	0.261	0.393	1.939	0.0749	0.0112	0.0929	1.032	4.639

which takes different coefficients for $u_k(r)$ and $v_k(r)$. After getting eigenvalues and wave functions for the 0-th order solutions, we use those wave functions to calculate p/m_Q corrections in order by order. Corrections are estimated by evaluating matrix elements of corresponding terms in the higher order Hamiltonian. Details of the prescription for carrying out numerical calculations and getting reliable solutions are described in [24].

In the work of [24], by fitting the smallest eigenvalues of the Hamiltonian with masses of $D(1867)$ and $D^*(2008)$ for $c\bar{q}$ ($q = u, d$), $D_s(1969)$ and $D_s^*(2112)$ for $c\bar{s}$, and $B(5279)$ and $B(5325)$ for $b\bar{q}$, a strong coupling constant α_s and other potential parameters, a and b , were determined. At that time following the paper [32], we started the search for a set of parameters with $b < 0$. Using those parameters obtained this way, other mass levels were calculated and compared with the experimental data for $D_{(s)}/B_{(s)}$ mesons. Light quark (u and d) mass was taken to be 10MeV as an input, being close to a value of current quark mass.

Here in this paper since we have now a plenty of the observed data, we follow a different way of analysis.

- 1) First, we determine the six parameters, i.e., a strong coupling constant α_s^c , potential parameters a and b , quark masses $m_{u,d}$ ⁴, m_s , and m_c by fitting calculations to the observed six D meson masses, i.e., those of two $\ell = 0$ ($0^-, 1^-$) states and four $\ell = 1$ ($0^+, 1^+, 1^+, 2^+$) states, and similarly six D_s meson masses, using the Minuit χ^2 analysis. The results are given in Table II. As written in the Introduction, we have found that the optimal value for b becomes positive contrary to the former paper [24], which affects all other parameter values as well as the calculated values, M_{calc} , M_0 , p_i , n_i , and c_i whose meanings are explained below.
- 2) Then, using the optimal parameters obtained this way except for a strong coupling, the bottom quark mass m_b and a strong coupling α_s^b which is assumed to be different from α_s^c for D/D_s mesons are determined by fitting the four observed B meson masses and two B_s meson masses. The most optimal values of parameters are presented in Table II at the first order calculation in p/m_Q .

The masses with the same value of k degenerate in the 0-th order for members of each spin doublet, are labeled as M_0 . c_i denotes the i -th order correction in p/m_Q expansion (in this paper $n = 1, 2$) and thus, the calculated heavy meson mass M_{calc} is given by the sum of M_0 and corrections up to the n -th order,

$$M_{\text{calc}} = M_0 + \sum_{i=1}^n (p_i + n_i), \quad (21)$$

where p_i includes the contributions stemming from only upper components of a heavy quark, n_i from lower, and $c_i = p_i + n_i$ in each order. Here one should notice that $M_0 = m_Q + E_0$. The degeneracy in k or for the states in the same spin doublet is resolved by taking account of higher order corrections in p/m_Q expansion. See Fig. 1.

In our formalism, each state is uniquely classified by two quantum numbers, k and j , and also approximately classified by the upper component of the wave function of Eq.(9) in terms of the conventional notation $^{2s+1}\ell_j$. In terms of this notation, the $j^P = 1^+$ state is a mixed state of 3P_1 and 1P_1 , while the $j^P = 2^-$ state is a mixed state of 3D_2 and 1D_2 . We approximately denote them with double quotations in Tables III–VII. The relations between them are given by [24]

$$\begin{pmatrix} |^3P_1''\rangle \\ |^1P_1''\rangle \end{pmatrix} = \frac{1}{\sqrt{3}} \begin{pmatrix} \sqrt{2} & 1 \\ -1 & \sqrt{2} \end{pmatrix} \begin{pmatrix} |^3P_1\rangle \\ |^1P_1\rangle \end{pmatrix}, \quad (22)$$

$$\begin{pmatrix} |^3D_2''\rangle \\ |^1D_2''\rangle \end{pmatrix} = \frac{1}{\sqrt{5}} \begin{pmatrix} \sqrt{3} & \sqrt{2} \\ -\sqrt{2} & \sqrt{3} \end{pmatrix} \begin{pmatrix} |^3D_2\rangle \\ |^1D_2\rangle \end{pmatrix}. \quad (23)$$

⁴ In the numerical analysis, we set $m_u = m_d$ in order to reduce the number of parameters.

TABLE III: D meson mass spectra (units are in MeV).

$^{2s+1}L_J(J^P)$	M_0	c_1/M_0	p_1/M_0	n_1/M_0	M_{calc}	M_{obs}
$^1S_0(0^-)$	1784	0.476×10^{-1}	0.374×10^{-1}	1.013×10^{-2}	1869	1867
$^3S_1(1^-)$		1.271×10^{-1}	1.266×10^{-1}	0.512×10^{-3}	2011	2008
$^3P_0(0^+)$	2067	1.046×10^{-1}	0.959×10^{-1}	0.874×10^{-2}	2283	2308
$^3P_1(1^+)$		1.713×10^{-1}	1.689×10^{-1}	2.444×10^{-3}	2421	2427
$^1P_1(1^+)$	2125	1.415×10^{-1}	1.410×10^{-1}	0.486×10^{-3}	2425	2420
$^3P_2(2^+)$		1.618×10^{-1}	1.617×10^{-1}	1.364×10^{-4}	2468	2460
$^3D_1(1^-)$	2322	1.894×10^{-1}	1.872×10^{-1}	2.228×10^{-3}	2762	—
$^3D_2(2^-)$		2.054×10^{-1}	2.052×10^{-1}	1.248×10^{-4}	2800	—

TABLE IV: D_s meson mass spectra (units are in MeV).

$^{2s+1}L_J(J^P)$	M_0	c_1/M_0	p_1/M_0	n_1/M_0	M_{calc}	M_{obs}
$^1S_0(0^-)$	1900	0.352×10^{-1}	0.270×10^{-1}	0.816×10^{-2}	1967	1969
$^3S_1(1^-)$		1.102×10^{-1}	1.098×10^{-1}	4.076×10^{-4}	2110	2112
$^3P_0(0^+)$	2095	1.101×10^{-1}	1.027×10^{-1}	0.740×10^{-2}	2325	2317
$^3P_1(1^+)$		1.779×10^{-1}	1.752×10^{-1}	2.620×10^{-3}	2467	2460
$^1P_1(1^+)$	2239	1.274×10^{-1}	1.270×10^{-1}	3.860×10^{-4}	2525	2535
$^3P_2(2^+)$		1.467×10^{-1}	1.466×10^{-1}	1.035×10^{-4}	2568	2572
$^3D_1(1^-)$	2342	2.032×10^{-1}	2.008×10^{-1}	2.382×10^{-3}	2817	—
$^3D_2(2^-)$		2.196×10^{-1}	2.195×10^{-1}	0.989×10^{-4}	2856	—

A. D and D_s

Calculated and optimally fitted values M_{calc} of D and D_s meson masses with the first order corrections are presented in Tables III and IV, respectively, together with experimental data M_{obs} . Those mass spectra of D and D_s mesons are also plotted in Figs. 2 and 3 and Figure 3 also includes DK/D^*K threshold lines. As one can see easily, our calculated masses M_{calc} are in good agreement with each observed value M_{obs} , and the discrepancies are less than 1%. Especially it is remarkable that newly observed levels $^3P_0(0^+)$ and $^3P_1(1^+)$ of both D and D_s mesons are reproduced well, where the masses of D_{sJ} mesons are below DK/D^*K thresholds. These levels of D and D_s mesons cannot be reproduced by any other quark potential models which predict about 100 ~ 200 MeV higher than our values even though they also succeed in reproducing other levels, i.e., $^1S_0(0^-)$, $^3S_1(1^-)$, $^1P_1(1^+)$, and $^3P_2(2^+)$.⁵ This result gives us a great confidence that our framework may give a good solution to the narrow D_{sJ} puzzle.

The first order corrections for D and D_s mesons are moderately large, whose amount is an order of 10%. Since the charm quark mass in our fitting is rather small, second order corrections might be necessary to get more reliable results. However, we have found that they are only an order of 1% or less and thus have neglected them in this work. To demonstrate this smallness, we also present the fit for D meson with the second order corrections in Table V as a typical example. When one looks at this Table, one notices that the fit with the experiments does not increase accuracy compared with Table III of the first order. Hence from here on in our computation we use only the first order calculations to obtain any levels including radial excitations which we present in a separate paper [35].

B. B and B_s

For B and B_s mesons, unfortunately there are only a few data available although newly discovered states ($B_1(5720)$, $B_2^*(5745)$, and $B_{s2}^*(5839)$) are reported recently by D0 and CDF.[5] Calculated masses of B and B_s mesons are given in Tables VI and VII, respectively. We have used a new value of a strong coupling α_s^b given by Table II for calculating B and B_s mesons different from α_s^c for D meson in Table II. Newly discovered levels are well reproduced by our model,

⁵ See Table XIV of [23].

too, as shown in these Tables. We predict the mass of several excited states which are not yet observed. The mass spectra of B and B_s mesons are also plotted in Figs. 4 and 5 and Figure 5 also includes BK/B^*K threshold lines. Among these levels, predicted masses of 0^+ and 1^+ states for B_s mesons are below BK/B^*K threshold the same as the case for D_{sJ} mesons. Therefore, their decay modes are kinematically forbidden, and the dominant decay modes are the pionic decay which is the same results obtained in [25] although values of B_s mesons are slightly different from each other:

$$B_s(0^+) \rightarrow B_s(0^-) + \pi, \quad (24a)$$

$$B_s(1^+) \rightarrow B_s(1^-) + \pi. \quad (24b)$$

The decay widths of these states are expected to be narrow the same as D_{sJ} meson cases, since those decay modes of Eq.(24) violate the isospin invariance. These higher states might be observed in Tevatron/LHC experiments or even in D0 and/or CDF of FNAL in the near future by analyzing the above decay modes. This is because D0 and CDF have recently announced the discovery of $B_1(5720) = B(^1P_1(1^+))$, $B_2^*(5745) = B(^3P_2(2^+))$, and $B_{s2}^*(5839) = B_s(^3P_2(2^+))$, which have narrow decay width with the same decay products as in Eq.(24) since only the D-wave decay channel is possible.[5] Even though these mass values are reproduced by other models [9, 21, 36, 37], but again those models predict larger mass values for 0^+ and 1^+ states of B_s mesons than ours, the same situation as in D_{sJ} so that these states have broad decay width contrary to our prediction. We hope that our prediction could be tested in the forthcoming experiments which distinguish ours from other models.

IV. CONCLUSION AND DISCUSSION

Given recent new data and expanding a parameter space with positive b , we have reanalyzed the mass spectra of heavy mesons, D , D_s , B and B_s with our semirelativistic quark potential model, in which an effective Hamiltonian has both a heavy quark symmetry and a chiral symmetry in a certain limit of parameters. Hamiltonian, wave function, and eigenvalue are all consistently expanded in p/m_Q , and a light antiquark is treated as four-spinor Dirac particle while a heavy quark is nonrelativistically reduced using the FWT transformation. Calculated masses of not only newly observed excited ($\ell = 1$) states, $D_{s0}(2317)$, $D_{s1}(2460)$, $D_0^*(2308)$ and $D_1'(2427)$ which are not explained by conventional quark potential models, but also already established ($\ell = 0$ and $\ell = 1$) states of D and D_s mesons are simultaneously reproduced in good agreement with experimental data within one percent of accuracy. Now we also have plausible values of masses for 3D_1 and 3D_2 which we failed to give in the former paper [24]. This has been achieved by finding the most optimal parameter set with positive parameter b .

These results suggest that our model could be a good solution to the narrow D_{sJ} puzzle, which is considered to be still challenging and people are still looking for exotic state possibilities. We would say that the quark potential model still remains powerful enough to understand the heavy meson spectroscopy. The important thing in a heavy-light system is to expand all quantities in the model in p/m_Q and set equations consistently order by order and not to neglect the lower components of a heavy quark, which has not been taken into account by other quark potential models. We have also predicted the masses of higher excited states of B and B_s mesons which are not yet observed, at the first order in p/m_b . Among these, the newly discovered states by D0 and CDF are also reproduced by our model, too. There are other models to reproduce these states [9, 21, 36, 37] but they cannot predict 0^+ and 1^+ states less than BK/B^*K threshold. This is the main difference of the results when comparing our model with others. Other models for heavy mesons cannot yield the masses of $^3P_0(0^+)$ and $^3P_1(1^+)$ states of D_s and B_s mesons much lower than $^1P_1(1^+)$ and $^3P_2(2^+)$ states like ours.

TABLE V: Second order D meson mass spectra (units are in MeV).

$^{2s+1}\ell_j (j^P)$	M_0	c_1/M_0	c_2/M_0	M_{calc}	M_{obs}
$^1S_0(0^-)$	1783	0.678×10^{-1}	-2.091×10^{-2}	1867	1867
$^3S_1(1^-)$		1.245×10^{-1}	2.507×10^{-3}	2009	2008
$^3P_0(0^+)$	1935	1.694×10^{-1}	1.567×10^{-2}	2293	2308
$^3P_1(1^+)$		2.132×10^{-1}	1.665×10^{-3}	2350	2427
$^1P_1(1^+)$	2045	1.412×10^{-1}	2.499×10^{-3}	2432	2420
$^3P_2(2^+)$		1.616×10^{-1}	-1.003×10^{-3}	2448	2460
$^3D_1(1^-)$	2127	1.863×10^{-1}	2.109×10^{-4}	2803	—
$^3D_2(2^-)$		2.021×10^{-1}	-1.409×10^{-3}	2726	—

TABLE VI: B meson mass spectra (units are in MeV).

$^{2s+1}L_J(J^P)$	M_0	c_1/M_0	p_1/M_0	n_1/M_0	M_{calc}	M_{obs}
$^1S_0(0^-)$	5277	-0.161×10^{-2}	-3.795×10^{-3}	2.187×10^{-3}	5270	5279
$^3S_1(1^-)$		0.981×10^{-2}	9.706×10^{-3}	1.107×10^{-4}	5329	5325
$^3P_0(0^+)$	5570	0.401×10^{-2}	1.937×10^{-3}	2.072×10^{-3}	5592	—
$^3P_1(1^+)$		1.412×10^{-2}	1.400×10^{-2}	1.227×10^{-4}	5649	—
$^1P_1(1^+)$	5660	1.069×10^{-2}	1.066×10^{-2}	0.289×10^{-4}	5720	5720
$^3P_2(2^+)$		1.364×10^{-2}	1.364×10^{-2}	2.120×10^{-7}	5737	5745
$^3D_1(1^-)$	5736	2.203×10^{-1}	2.202×10^{-1}	4.583×10^{-5}	6999	—
$^3D_2(2^-)$		1.430×10^{-1}	1.430×10^{-1}	2.092×10^{-7}	6556	—

TABLE VII: B_s meson mass spectra (units are in MeV).

$^{2s+1}L_J(J^P)$	M_0	c_1/M_0	p_1/M_0	n_1/M_0	M_{calc}	M_{obs}
$^1S_0(0^-)$	5394	-0.302×10^{-2}	-0.485×10^{-2}	0.183×10^{-2}	5378	5369
$^3S_1(1^-)$		0.853×10^{-2}	0.844×10^{-2}	9.249×10^{-5}	5440	—
$^3P_0(0^+)$	5598	0.350×10^{-2}	0.173×10^{-2}	0.177×10^{-2}	5617	—
$^3P_1(1^+)$		1.498×10^{-2}	1.444×10^{-2}	5.396×10^{-4}	5682	—
$^1P_1(1^+)$	5775	0.978×10^{-2}	0.969×10^{-2}	8.257×10^{-5}	5831	—
$^3P_2(2^+)$		1.263×10^{-2}	1.261×10^{-2}	2.014×10^{-5}	5847	5839
$^3D_1(1^-)$	5875	2.949×10^{-2}	2.898×10^{-2}	5.104×10^{-4}	6048	—
$^3D_2(2^-)$		0.564×10^{-2}	0.562×10^{-2}	1.980×10^{-5}	5908	—

Some comments are in order.

(i) As shown in Tables III–VII, the first order corrections to the 0-th order mass, c_1/M_0 , are roughly an order of 10% for D and D_s mesons, while 1% for B and B_s mesons. Therefore, p/m_Q expansion works much better for B and B_s mesons. This is quite natural, since the bottom quark mass is heavier than the charm quark mass. In addition, it is interesting that the first order correction becomes larger for higher excited states. It means that the relativistic correction must be more important in higher excited states, since the higher order corrections in p/m_Q expansion represent the relativistic effects of a heavy quark. This is consistent with a naive picture that the inner motion of light as well as heavy quarks may be larger in excited states, and therefore the effects of lower components of a heavy quark may become much more important to be included.

(ii) The mass difference between members of a spin doublet satisfies the following relation (given by Eq.(53) of [24]):

$$m_c(M_{D^*} - M_D) = m_b(M_{B^*} - M_B), \quad (25)$$

$$m_c(M_{D_s^*} - M_{D_s}) = m_b(M_{B_s^*} - M_{B_s}), \quad (26)$$

in the first order of calculation. This is because each of two states having the same value of the quantum number k has an equal mass at the 0-th order and hence the mass is degenerate at this stage as shown in Fig. 1. The splitting occurs by including the first order (p/m_Q) corrections originated from H_1 in Eq.(4), which is proportional to $f^k(m_q)/m_Q$ with the same functional form $f^k(m_q)$ for two states, and therefore Eqs.(25, 26) are exactly satisfied in our model. Similar relations hold for higher spin states with the same value of k .

(iii) As shown in Table II, the light quark masses $m_{u,d}$ and m_s are considerably small, $m_{u,d} \sim 10$ MeV and $m_s \sim 90$ MeV, compared with the constituent quark masses which are adopted in conventional potential models [7]. It should be noted that these values are not input but the outcome of this analysis, though in the previous analysis [24], $m_{u,d} = 10$ MeV is taken as an input. This result is consistent with those obtained in [38, 39, 40].

This situation might be described as follows. A light quark is moving around a heavy quark in a heavy meson wearing small gluon clouds with large momenta in our treatment of quarks so that light quarks have so-called current quark mass. This holds only in heavy-light systems and in the other systems, e.g., quarkonium and ordinary hadrons, we should use quarks wearing more gluon clouds, i.e., constituent quarks as adopted in usual quark potential models. There may be another interpretation bridging our model with light mass with a constituent quark model.

(iv) Repeatedly saying, the main difference between other conventional quark potential models and ours is that our model systematically expands interaction terms in p/m_Q , takes into account lower component contributions of




FIG. 2: Plot of D meson masses in GeV. Solid bars and white circles represent calculated and observed masses, respectively. Specific values are given in Table III.




FIG. 3: Plot of D_s meson masses. The legend is as for Fig. 2. The dashed lines show DK and D^*K thresholds, respectively. Specific values are given in Table IV.

heavy quarks in the intermediate states when calculating higher order terms, and adopt very light antiquark mass. With regard to $1/m_Q$ expansion, for instance, the model of [7, 8] has kinetic terms, $\sqrt{p^2 + m^2}$, while they adopted nonrelativistic interaction terms which are symmetric in Q and $\bar{q}$ or in m_Q and $m_{\bar{q}}$. The model of [9] somehow manipulated to hold heavy quark symmetry and expanded interactions in $1/m_Q$ but adopted constituent masses for light quarks, m_u , m_d , and m_s . Even so, their numerical results give about a hundred MeV larger values for D_{sJ} particles. Both models have not taken into account the lower components of a heavy quark which naturally appear in our semirelativistic potential model.

(v) We have been successful in reproducing experimental data for both heavy mesons D/D_s and B/B_s . However we have also left unresolved problems in our semirelativistic model. When one looks at Table II, one notices that values of a strong coupling used for D/D_s and B/B_s are largely different from each other. Especially that for D/D_s is too small to be a strong coupling. Also we have not used a running strong coupling although most of other quark potential models take into account this effect in some way.

Acknowledgments

One of the authors (K.S.) would like to thank S. Yokkaichi for helpful discussions about numerical calculations.




FIG. 4: Plot of B meson masses. The legend is as for Fig. 2. Specific values are given in Table VI. Calculated mass values for 3D_1 and 3D_2 are too large to put them in the same figure with others.




FIG. 5: Plot of B_s meson masses. The legend is as for Fig. 2. The dashed lines show BK and B^*K thresholds, respectively. Specific values are given in Table VII.

APPENDIX A: SCHRÖDINGER EQUATION WITH AN EFFECTIVE HAMILTONIAN

The original Schrödinger equation is actually modified by the FWT transformation in our framework in order to make non-relativistic reduction of a heavy quark. Now we recall the modified Schrödinger equation of Eq. (3),

$$(H_{\text{FWT}} - m_Q) \otimes \psi_{\text{FWT}} = \tilde{E} \psi_{\text{FWT}}, \quad (\text{A1})$$

The Hamiltonian and the wave function are transformed by the following FWT and charge conjugate operator on the heavy quark sector:

$$H_{\text{FWT}} = U_c U_{\text{FWT}}(p'_Q) H U_{\text{FWT}}^{-1}(p_Q) U_c^{-1}, \quad (\text{A2})$$

$$\psi_{\text{FWT}} = U_c U_{\text{FWT}}(p_Q) \psi, \quad (\text{A3})$$

where given are

$$U_{\text{FWT}}(p) = \exp\left(W(p) \vec{\gamma}_Q \cdot \vec{p}\right) = \cos W(p) + \vec{\gamma}_Q \cdot \vec{p} \sin W(p), \quad (\text{A4})$$

$$U_c = i\gamma_Q^0 \gamma_Q^2 = -U_c^{-1}, \quad (\text{A5})$$

and some kinematical variables are defined as

$$\vec{\tilde{p}} = \frac{\vec{p}}{p}, \quad \tan W(p) = \frac{p}{m_Q + E}, \quad E = \sqrt{\vec{p}^2 + m_Q^2}, \quad (\text{A6})$$

$$\vec{p} = \vec{p}_q = -\vec{p}_Q, \quad \vec{p}' = \vec{p}'_q = -\vec{p}'_Q, \quad \vec{q} = \vec{p}' - \vec{p}. \quad (\text{A7})$$

Here we are considering a scattering of a heavy quark Q and a light antiquark $\bar{q}$. Hence, $\vec{p}_i$, $\vec{p}'_i$, and $\vec{q}$ given above represent momenta of incoming, outgoing quarks, and gluon which is exchanged between quarks, respectively. Note that the argument of the FWT transformation U_{FWT} operating on a Hamiltonian from left is different from the right-operating one, since an outgoing momentum $\vec{p}'_Q$ is different from an incoming one $\vec{p}_Q$. However, in our study we work in a configuration space in which momenta are nothing but the derivative operators. When we write them differently, for instance as $\vec{p}_Q$ and $\vec{p}'_Q$, their expressions are reminders of their momentum representation. Therefore, although the arguments of U_{FWT} and U_{FWT}^{-1} look different, $\vec{p}_Q$ and $\vec{p}'_Q$ are expressed by the same derivative operator $-i\vec{\nabla}$. The momentum transfer $\vec{q}$ operates only on potentials and provides nonvanishing results.

The charge conjugation operator U_c is introduced to make the wave function $U_c\psi$ a true bi-spinor, i.e., gamma matrices of a light antiquark are multiplied from left while those of a heavy quark from right, which is expressed by using a notation $\otimes$.

Then, the Hamiltonian in the modified Schrödinger equation given by Eq. (3) is expanded in powers of p/m_Q :

$$H_{\text{FWT}} - m_Q = H_{-1} + H_0 + H_1 + H_2 + \cdots. \quad (\text{A8})$$

H_i stands for the i -th order expanded Hamiltonian whose explicit forms are given by

$$H_{-1} = -(1 + \beta_Q)m_Q, \quad (\text{A9a})$$

$$H_0 = \vec{\alpha}_q \cdot \vec{p} + \beta_q m_q - \beta_q \beta_Q S + \left\{ 1 + \frac{1}{2} [\vec{\alpha}_q \cdot \vec{\alpha}_Q + (\vec{\alpha}_q \cdot \vec{n})(\vec{\alpha}_Q \cdot \vec{n})] \right\} V, \quad (\text{A9b})$$

$$\begin{aligned} H_1 = & -\frac{1}{2m_Q} \beta_Q \vec{p}^2 + \frac{1}{m_Q} \beta_q \vec{\alpha}_Q \cdot \left(\vec{p} + \frac{1}{2} \vec{q} \right) S + \frac{1}{2m_Q} \vec{\gamma}_Q \cdot \vec{q} V \\ & - \frac{1}{2m_Q} \left[\beta_Q \left(\vec{p} + \frac{1}{2} \vec{q} \right) + i\vec{q} \times \beta_Q \vec{\Sigma}_Q \right] \cdot [\vec{\alpha}_q + (\vec{\alpha}_q \cdot \vec{n})\vec{n}] V, \end{aligned} \quad (\text{A9c})$$

$$\begin{aligned} H_2 = & \frac{1}{2m_Q^2} \beta_q \beta_Q \left(\vec{p} + \frac{1}{2} \vec{q} \right)^2 S - \frac{i}{4m_Q^2} \vec{q} \times \vec{p} \cdot \beta_q \beta_Q \vec{\Sigma}_Q S - \frac{1}{8m_Q^2} \vec{q}^2 V - \frac{i}{4m_Q^2} \vec{q} \times \vec{p} \cdot \vec{\Sigma}_Q V \\ & - \frac{1}{8m_Q^2} \{ (\vec{p} + \vec{q})(\vec{\alpha}_Q \cdot \vec{p}) + \vec{p} [\vec{\alpha}_Q \cdot (\vec{p} + \vec{q})] + i\vec{q} \times \vec{p} \gamma_Q^5 \} \cdot [\vec{\alpha}_q + (\vec{\alpha}_q \cdot \vec{n})\vec{n}] V, \end{aligned} \quad (\text{A9d})$$

$\vdots$

More details of matrix elements of each order in $1/m_Q$ and properties of wave functions with several operators are given and evaluated in Ref. [24].

Eq.(A9a) means that the lowest order equation gives $(1 + \beta_Q)\psi = 0$, which implies that a heavy quark is regarded as a static source of color so that projection operator becomes $(1 \pm \beta_Q)/2$, the one at rest system. That is, the solution given by Eq.(8) includes only upper or positive components with regard to a heavy quark. In this sense, upper/lower component of a heavy quark has the same meaning as positive/negative energy component.

APPENDIX B: BLOCH METHOD

This Appendix is based on the paper [30], by which we explain Bloch's method [31] for a degenerate system. We will see the Foldy-Wouthuysen-Tani [29] transformation is automatically carried out and will obtain the systematic expansion of the heavy-light system in $1/m_Q$. To distinguish this formulation from the one we have used, we use different notations in this Appendix.

We expand the eigenvalues of the Hamiltonian with respect to the inverse of m_Q . First we divide the Hamiltonian as $\mathcal{H} = \beta_Q m_Q + \mathcal{V}$, and we treat $\mathcal{V}$ as a perturbation, where actually $\mathcal{H} = H$. The eigenvalues of the unperturbed Hamiltonian are simply $\pm m_Q$, and the eigenstates of $\mathcal{H}$ are almost degenerate. Then we apply the perturbation method of Bloch [31]. In general, let us consider the Hamiltonian $\mathcal{H} = H_0 + \mathcal{V}$. We denote the projection operator to the subspace M with an eigenvalue E of H_0 as P , and the projection operator to the subspace $\mathcal{M}$ with eigenvalues $\{\mathcal{E}_j\}$ of $\mathcal{H}$ as $\mathcal{P}$, where all $\mathcal{E}_j$ converge to E in the vanishing limit of the interaction $\mathcal{V}$. Further we choose appropriate vectors $\{\psi_j\}$ in the subspace M such that their projections on $\mathcal{M}$ become the eigenvectors of $\mathcal{H}$, that is, $\mathcal{H}\mathcal{P}\psi_j = \mathcal{E}_j\mathcal{P}\psi_j$. If we define an operator $\mathcal{U}$ by $\mathcal{P}\mathcal{P} = \mathcal{U}\mathcal{P}\mathcal{P}$, then we get the following eigenvalue equation,

$$\mathcal{H}_{eff}(\mathcal{P}\mathcal{P}\psi_j) = \mathcal{E}_j(\mathcal{P}\mathcal{P}\psi_j), \quad (\text{B1})$$

where the effective Hamiltonian $\mathcal{H}_{eff}$ is defined by $\mathcal{H}_{eff} = \mathcal{P}\mathcal{H}\mathcal{U}$. Here we must notice that this effective Hamiltonian is not hermitian. The eigenfunction of $\mathcal{H}$, $\mathcal{P}\psi_j$, is obtained from the eigenfunction of $\mathcal{H}_{eff}$ given by $\phi_j = \mathcal{P}\mathcal{P}\psi_j$, by multiplying $\mathcal{U}$ from left, that is,

$$\mathcal{U}\phi_j = \mathcal{U}\mathcal{P}\mathcal{P}\psi_j = \mathcal{P}\psi_j. \quad (\text{B2})$$

Note that $\mathcal{P}\psi_j = \phi_j$. Again we must notice that $\mathcal{U}$ is not unitary, therefore, even if ϕ_j is normalized, $\mathcal{U}\phi_j$ is not.

$\mathcal{P}$ and $\mathcal{U}$ are formally expanded as follows.

$$\mathcal{P} = P - \sum_{m=1}^{\infty} \sum_{\{k_i\}} S_{k_1} \mathcal{V} S_{k_2} \cdots S_{k_m} \mathcal{V} S_{k_{m+1}}, \quad (\text{B3})$$

where S_k 's are defined by

$$S_0 = -P \quad \text{and} \quad S_k = \frac{Q}{a^k} \quad (k = 1, 2, \cdots) \quad \text{with} \quad Q = 1 - P, \quad a = E - H_0, \quad (\text{B4})$$

and the sum over $\{k_i\}$ extends over non-negative integers satisfying the condition $\sum_{i=1}^{m+1} k_i = m$.

$$\mathcal{U} = \sum_{m=1}^{\infty} \sum'_{\{k_i\}} S_{k_1} \mathcal{V} S_{k_2} \cdots S_{k_m} \mathcal{V} P, \quad (\text{B5})$$

where the sum over $\{k_i\}$ extends over non-negative integers satisfying the conditions $\sum_{i=1}^m k_i = m$ and $\sum_{i=1}^p k_i \geq p$ ($p = 1, 2, \cdots m-1$). Making use of the expansion formula for $\mathcal{U}$, we obtain the effective Hamiltonian including the second order corrections in $1/a \sim 1/m_Q$ as follows:

$$\mathcal{H}_{eff} = EP + P\mathcal{V}P + P\mathcal{V}\frac{Q}{a}\mathcal{V}P + P\mathcal{V}\frac{Q}{a}\mathcal{V}\frac{Q}{a}\mathcal{V}P - P\mathcal{V}\frac{Q}{a^2}\mathcal{V}P\mathcal{V}P + \cdots. \quad (\text{B6})$$

where $P = (1 + \beta_Q)/2$ and $Q = (1 - \beta_Q)/2$.

-
- [1] BaBar Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **90**, 242001 (2003).
 - [2] CLEO Collaboration, D. Besson *et al.*, Phys. Rev. D **68**, 032002 (2003).
 - [3] Belle Collaboration, P. Krokovny *et al.*, Phys. Rev. Lett. **91**, 262002 (2003); Y. Mikami *et al.*, Phys. Rev. Lett. **92**, 012002 (2004).
 - [4] Belle Collaboration, K. Abe *et al.*, Phys. Rev. D **69**, 112002 (2004).
 - [5] CDF Collaboration, CDF note 7938 (2005); D0 Collaboration, D0notes 5026-CONF, 5027-CONF (2006); I. Kravchenko, hep-ex/0605076.
 - [6] T. Barnes, F. E. Close, and H. J. Lipkin, Phys. Rev. D **68**, 054006 (2003).
 - [7] S. Godfrey and N. Isgur, Phys. Rev. D **32**, 189 (1985).
 - [8] S. Godfrey and R. Kokoski, Phys. Rev. D **43**, 1679 (1991).
 - [9] D. Ebert, V.O. Galkin, and R.N. Faustov, Phys. Rev. D **57**, 5663 (1998).
 - [10] S. N. Mukherjee, R. Nag, S. Sanyal, T. Morii, J. Morishita, and M. Tsuge, Phys. Rept. **231**, 201 (1993).
 - [11] W. A. Bardeen, E. J. Eichten, and C. T. Hill, Phys. Rev. D **68**, 054024 (2003).
 - [12] W. A. Bardeen and C. T. Hill, Phys. Rev. D **49**, 409 (1994); M. A. Nowak, M. Rho, and I. Zahed, Phys. Rev. D **48**, 4370 (1993); D. Ebert, T. Feldmann, and H. Reinhardt, Phys. Lett. B **388**, 154 (1996); A. Deandrea, N. Di Bartolomeo, R. Gatto, G. Nardulli, and A. D. Polosa, Phys. Rev. D **58**, 034004 (1998); A. Hiorth and J. O. Eeg, Phys. Rev. D **66**, 074001 (2002); M. Harada, M. Rho, and C. Sasaki, Phys. Rev. D **70**, 074002 (2004).

- [13] A. Hayashigaki and K. Terasaki, hep-ph/0411285.
- [14] A. P. Szczepaniak, Phys. Lett. B **567**, 23 (2003).
- [15] T. E. Browder, S. Pakvasa, and A. A. Petrov, Phys. Lett. B **578**, 365 (2004).
- [16] K. Terasaki, Phys. Rev. D **68**, 011501(R) (2003); K. Terasaki and B. H. J. McKellar, hep-ph/0501188.
- [17] V. Dmitrašinović, Phys. Rev. Lett. **94**, 162002 (2005).
- [18] G. S. Bali, Phys. Rev. D **68**, 071501(R) (2003); UKQCD Collaboration, A. Dougall, R. D. Kenway, C. M. Maynard, and C. McNeile, Phys. Lett. B **569**, 41 (2003).
- [19] C. Habe, K. Iwata, M. Hirano, T. Murota, and D. Tsuruda, Prog. Theor. Phys. **77**, 917 (1987).
- [20] A. Gara, B. Durand, and L. Durand, Phys. Rev. D **40**, 843 (1989).
- [21] M. Di Piero and E. Eichten, Phys. Rev. D **64**, 114004 (2001); see also J.L. Goity and W. Roberts, Phys. Rev. D **60**, 034001 (1999) and J. Zeng, J. W. Van Orden, and W. Roberts, Phys. Rev. D **52**, 5229 (1993).
- [22] E. van Beveren and G. Rupp, Phys. Rev. Lett. **97**, 202001 (2006); Mod. Phys. Lett. A19, 1949 (2004); Eur. Phys. J. C **32**, 493 (2004); Phys. Rev. Lett. **91**, 012003 (2003).
- [23] E. S. Swanson, Phys. Rept. **429**, 243 (2006), hep-ph/0601110.
- [24] T. Matsuki and T. Morii, Phys. Rev. D **56**, 5646 (1997); See also T. Matsuki, Mod. Phys. Lett. A **11**, 257 (1996), T. Matsuki and T. Morii, Aust. J. Phys. **50**, 163 (1997), T. Matsuki, T. Morii, and K. Seo, Trends in Applied Spectroscopy, **4**, 127 (2002); T. Matsuki, T. Morii, and K. Sudoh, AIP Conference Proceedings, Vol. 814, p.533 (2005), "XI International Conference on Hadron Spectroscopy" (Hadron05), held at Rio De Janeiro, Brazil 21-26 Aug. 2005, hep-ph/0510269 (2005).
- [25] T. Matsuki, K. Mawatari, T. Morii, and K. Sudoh, Phys. Lett. B **606**, 329 (2005).
- [26] E. Fermi and C. N. Yang, Phys. Rev. **34**, 553 (1949).
- [27] For example, H. Bethe and E. E. Salpeter, *Handbuch der Physik*, edited by H. von S. Flüge, Band XXXV ATOMS I (Springer, Berlin, 1955), p. 256.
- [28] H. Miyazawa and K. Tanaka, Prog. Theor. Phys. **87**, 1457 (1992).
- [29] L. L. Foldy and S. A. Wouthuysen, Phys. Rev. **78**, 29 (1950); S. Tani, Prog. Theor. Phys. **6**, 267 (1951).
- [30] T. Matsuki and K. Seo, Bulletin of Gifu City Women's College **46**, 7 (1997).
- [31] C. Bloch, Nucl. Phys. **6**, 329 (1958); A. Messiah, *Quantum Mechanics* (North-Holland, Amsterdam), p.718 (1962).
- [32] J. Morishita, M. Kawaguchi, and T. Morii, Phys. Lett. B **185**, 139 (1987); Phys. Rev. D **37**, 159 (1988).
- [33] M. E. Rose, *Relativistic Electron Theory*, John Wiley & Sons, 1961; J. J. Sakurai, *Advanced Quantum Mechanics*, Addison-Wesley, 1967.
- [34] T. Matsuki, K. Mawatari, T. Morii, and K. Sudoh, hep-ph/0408326.
- [35] T. Matsuki, T. Morii, and K. Sudoh, a talk given by T. Matsuki at the IVth International Conference on Quarks and Nuclear Physics (QNP06), held at Madrid, Spain 5-10 June 2006, hep-ph/0610186.
- [36] E. Eichten, C. Hill, and C. Quigg, Phys. Rev. Lett. **71**, 4116 (1993).
- [37] A. Hiorth Örsland and H. Högaasen, Eur. Phys. J. C9, 503 (1999).
- [38] H.D. Politzer, Nucl. Phys. B **117**, 397 (1976).
- [39] T. Matsuki and N. Yamamoto, Phys. Rev. D **22**, 2433 (1980).
- [40] M. Kaburagi, M. Kawaguchi, T. Morii, T. Kitazoe, and J. Morishita, Z. Phys. C **9**, 213 (1981).